

Hemocyanin Suggests a Close Relationship of Remipedia and Hexapoda

Beyhan Ertas,* Björn M. von Reumont,† Johann-Wolfgang Wägele,† Bernhard Misof,* and Thorsten Burmester*

*Biozentrum Grindel und Zoologisches Museum, Universität Hamburg, Hamburg, Germany; and †Zoologisches Forschungsmuseum Alexander Koenig, Bonn, Germany

The Remipedia are enigmatic crustaceans from anchialine cave systems, first described only 30 years ago, whose phylogenetic affinities are as yet unresolved. Here we report the sequence of hemocyanin from *Speleonectes tulumensis* Yager, 1987 (Remipedia, Speleonectidae). This is the first proof of the presence of this type of respiratory protein in a crustacean taxon other than Malacostraca. *Speleonectes tulumensis* hemocyanin consists of multiple distinct (at least three) subunits (StuHc1–3; Hc, hemocyanin). Surprisingly, the sequences are most similar to hexapod hemocyanins. Phylogenetic analyses showed that the *S. tulumensis* hemocyanin subunits StuHc1 and StuHc3 associate with the type 1 hexapod hemocyanin subunits, whereas StuHc2 associates with the type 2 subunits of hexapods. Together, remipede and hexapod hemocyanins are in the sister-group position to the hemocyanins of malacostracan crustaceans. Hemocyanins provide no indication of a close relationship of Myriapoda and Hexapoda but support Pancrustacea (Crustacea + Hexapoda). Our results also suggest that Crustacea are paraphyletic and that Hexapoda may have evolved from a Remipedia-like ancestor. Thus, Remipedia occupy a key position for the understanding of the evolution of hexapods, which are and have been one of the world's most speciose lineage of animals.

Introduction

In 1979, Yager discovered in an anchialine cave in the Bahamas the first specimens of Remipedia, which represented a new class of crustaceans with currently about 20 described extant species (Yager 1981; Koenemann, Schram, Hönemann, and Iliffe 2007). Living remipedes harbor a number of unique features, which include the loss of eyes, biramous antennulae, three pairs of postmandibular mouthparts adapted to a predatory feeding mode and to grooming, and the lack of tagmatization of the trunk (Schram and Lewis 1989; Koenemann, Schram, Iliffe, et al. 2007; van der Ham and Felgenhauer 2008). The phylogenetic affinities of Remipedia are still controversial. On the basis of the homonomous segmentation of their trunk, which supposedly represents the ancestral ground pattern in the evolution of Arthropoda, it has been suggested that Remipedia occupy a basal position within Crustacea or even Mandibulata (Schram 1986; Wills 1997; Giribet et al. 2001). Molecular phylogenetic analyses of ribosomal RNA and mitochondrial genomes, as well as limb morphology, have indicated an association of Remipedia with various crustacean classes assigned to the “Maxillopoda” (Ito 1989; Spears and Abele 1997; Lim and Hwang 2006). However, these results may be due to long branch attraction phenomena (Spears and Abele 1997; Telford et al. 2008; von Reumont et al. 2009). In fact, the pattern of mitochondrial gene arrangement convincingly excluded Remipedia from the maxillopod assemblage, which in that study comprised Branchiura, Cephalocarida, Cirripedia, and Pentastomida (Lavrov et al. 2004). Other molecular phylogenetic analyses have found Remipedia at various positions within Crustacea, albeit with poor support (Regier and Shultz 2001; Regier et al. 2005; Hassanin 2006; Carapelli et al. 2007). A recent study of arthropod brain morphology suggested that Remipedia belong to a monophylum of Malacostraca and Hexapoda within paraphyletic crustaceans

(Fanenbruck et al. 2004; Fanenbruck and Harzsch 2005). The discovery of free-living lecithotrophic remipede larvae and analysis of early larval development of Remipedia tentatively support their assumed relationship to malacostracan crustaceans (Koenemann, Schram, Bloechl, et al. 2007; Koenemann et al. 2009).

Oxygen transport in the hemolymph of various arthropod and mollusk taxa is facilitated by copper proteins referred to as hemocyanins (Markl and Decker 1992; van Holde and Miller 1995; Burmester 2002). Arthropod and mollusk hemocyanins are not related but evolved independently from distinct copper-containing enzymes (Burmester 2001, 2002). Hemocyanins of Arthropoda are large hexameric or oligohexameric proteins composed of similar or identical subunits in the size range of 75 kDa. Each subunit can bind to an O₂ molecule by the virtue of two copper ions that are coordinated by six histidine residues. Hemocyanins have been thoroughly studied in Chelicerata and Crustacea, occur in Myriapoda (Jaenicke et al. 1999; Kusche and Burmester 2001), and have recently been identified in Onychophora (Kusche et al. 2002) and Hexapoda (Hagner-Holler et al. 2004; Pick et al. 2008, 2009). Within Crustacea, hemocyanins have been thought to be confined to Malacostraca (Mangum 1985; Markl and Decker 1992). Hemocyanin sequences have been shown to be informative for the inference of phylogenies within Arthropoda (Burmester 2001; Kusche and Burmester 2001; Kusche et al. 2002, 2003). Thus, lineage-specific presence and phylogenetic analyses could be useful in assessing the position of Remipedia.

Here, we report the identification and molecular cloning of hemocyanin cDNA from the remipede *Speleonectes tulumensis* and show that molecular phylogenetic analyses of these hemocyanin sequences provide evidence for a close relationship of Remipedia and Hexapoda.

Materials and Methods

Sample Preparation

Speleonectes tulumensis was collected on Yucatan Peninsula, Mexico. Animals were cut into small pieces and the tissue was preserved in RNAlater (Qiagen, Hilden,

Key words: Crustacea, Hemocyanin, Hexapoda, Insecta, Pancrustacea, Remipedia.

E-mail: thorsten.burmester@uni-hamburg.de.

Mol. Biol. Evol. 26(12):2711–2718. 2009

doi:10.1093/molbev/msp186

Advance Access publication August 19, 2009

Germany). Total RNA and protein were extracted using TriFast (Peqlab, Erlangen, Germany) according to the manufacturer's instructions. The protein and RNA samples were immediately used or kept frozen at -20°C until use.

Cloning of Hemocyanin cDNA

Three micrograms of total RNA was converted into cDNA by Superscript III reverse transcriptase employing an oligo-dT primer according to the manufacturer's instructions (Invitrogen, Carlsbad, CA). The resulting cDNA was used for standard polymerase chain reaction (PCR), using degenerate primers (forward primer: 5'-ATGGAYTTYC CNTTYTGGTGGA-3'; reverse primer 5'-GTNGCGGTY TCRAARTGYTCCAT-3') that had been derived from conserved coding regions of arthropod hemocyanins (amino acid sequences: MDFPFWW and MEHFETAT) (Burmester 2001; Hagner-Holler et al. 2004; Pick et al. 2009). Fragments of the expected size were cloned into the pGem-T Easy/JM109 *Escherichia coli* system (Promega, Mannheim, Germany) and 14 independent clones were sequenced by a commercial service (Genterprise, Mainz, Germany). Rapid amplification of cDNA ends (RACE) experiments (5' and 3' RACE) were carried out by an RNA ligase-mediated rapid amplification method employing the GeneRacer Kit with SuperScript III reverse transcriptase. Sets of gene-specific primers were constructed according to the partial sequences (supplementary table 1, Supplementary Material online). The cDNA fragments were cloned and sequenced as described. Sequences were assembled using ContigExpress (Vector NTI Advance 10.3; Invitrogen) and GeneDoc 2.7 (Nicholas, Nicholas, and Deerfield 1997).

Western Blotting

Protein extracts were denatured in sample buffer (31.25 mM Tris-HCl, pH 6.8, 1% sodium dodecyl sulfate [SDS], 2.5% β -mercaptoethanol, and 5% glycerol) at 95°C for 5 min and loaded onto a 10% polyacrylamide gel. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was carried out according to standard procedures. Semidry electro-transfer of proteins onto nitrocellulose membranes (Hartenstein, Würzburg, Germany) was carried out for 2 h at 0.8 mA/cm^2 . Nonspecific binding sites were blocked for 1 h with 2% nonfat dry milk in tris-buffered saline (TBS) (10 mM Tris-HCl, pH 7.4; 140 mM NaCl). The membranes were incubated for 2 h at room temperature with polyclonal antibodies that had been raised against various insect and crustacean hemocyanins (anti-*Homarus americanus*, anti-*Panulirus interruptus*, anti-*Cancer pagurus*, and anti-*Thermobia domestica* hemocyanin antibodies) diluted 1:5,000 in 2% milk/TBS. The nitrocellulose filters were washed three times with TBS for 15 min and incubated for 1 h with a goat antirabbit antibody coupled with alkaline phosphatase (Dianova, Hamburg, Germany), diluted 1:10,000 in TBS. After a final washing step, detection was carried out with nitro-blue-tetrazolium chloride and 5-bromo-4-chloro-3-indolyl-phosphate as substrates.

Sequence and Phylogenetic Analyses

Tools provided with the ExPASy Molecular Biology Server of the Swiss Institute of Bioinformatics (<http://www.expasy.org>) were used for the analyses of DNA and amino acid sequences. Signal peptides were predicted using SignalP 1.1 (Nielsen et al. 1997). Deduced amino acid sequences of *S. tulumensis* hemocyanin were included in a previously published multiple alignment of arthropod hemocyanins, insect hexamerins, and crustacean pseudo-hemocyanins (Pick et al. 2009) employing MAFFT with the L-INS-i method and the BLOSUM 62 matrix (Katoh et al. 2005). A list of sequences used in this study is provided in supplementary table 1, Supplementary Material online.

After the exclusion of N- and C-terminal extensions, the final multiple sequence alignment contained 800 positions and 96 sequences. The appropriate model of amino acid sequence evolution (WAG + Gamma model; Whelan and Goldman 2001) was selected by ProtTest (Abascal et al. 2005) using the Akaike Information Criterion. Bayesian phylogenetic analysis was performed using MrBayes 3.1.2 (Huelsenbeck and Ronquist 2001). We assumed the WAG model with a gamma distribution of substitution rates. We used uninformative priors on all trees. Metropolis-coupled Markov chain Monte Carlo (MCMCMC) sampling was performed with one cold and three heated chains that were run for 8 million generations. Starting trees were random and trees were sampled every 100th generation. Two independent runs were performed in parallel and were continued until runs had converged (average standard deviations of split frequencies were stationary and lower than 0.005). The program Tracer 1.4 (<http://tree.bio.ed.ac.uk/software/tracer/>) was used to examine log-likelihood plots and MCMC summaries for all parameters. Posterior probabilities were estimated on the final 60,000 trees (burn-in = 20,000). RAxML 7.0.4, assuming the WAG evolutionary model with gamma distributions, was used for maximum likelihood (ML) analyses, and the resulting tree was tested by bootstrapping with 1,000 replicates. Tree-Puzzle 5.2 (Schmidt et al. 2002) was used to test alternative tree topologies. Hypothesis testing was performed by four methods: A Shimodaira-Hasegawa (SH) test (Shimodaira and Hasegawa 1999), a two-sided Kishino-Hasegawa (2sKH) test (Kishino and Hasegawa 1989), a one-sided KH (1sKH) test based on pairwise SH tests (Goldman et al. 2000), and an Expected Likelihood Weight (ELW) test (Strimmer and Rambaut 2002). SH, 1sKH, and ELW tests performed 1,000 resamplings using the REL method.

Results and Discussion

Identification of Hemocyanin in *S. tulumensis*

A set of degenerate oligonucleotide primers was designed according to conserved regions arthropod hemocyanin sequences (Burmester 2001; Pick et al. 2009). Reverse transcription-PCR using total RNA extracted from an adult *S. tulumensis* resulted in the amplification of fragments in the range of 550 bp. After cloning and sequencing, three distinct hemocyanin cDNA sequences were identified, which were termed StuHc1, StuHc2, and StuHc3, respectively (Hc, hemocyanin). The missing 5' and 3' regions of

protein or actually derives from the decapod host, *Carcinus maenas* (Herberts and de Frescheville 1981). Yager (1991) observed large crystal structures in remipede hemocytes that resemble those of the *Limulus polyphemus* hemocyanin. In a more recent report, van der Ham and Felgenhauer (2007) also speculated about the presence of a hemocyanin-like protein in *Speleonectes* sp. Here, we demonstrate that hemocyanin actually occurs in a remipede species, which is the first unambiguous report of such respiratory proteins in a non-malacostracan crustacean.

Remipedia dwell in a high-saline and oxygen-poor environment (usually <1mg/ml O₂; Koenemann et al. 2006) below the halocline interface between the seawater and the overlying well-oxygenated freshwater that is typical for anchialine cave systems (Pohlmann et al. 1997). Remipedia have adapted to this hypoxic environment, and a respiratory protein that augments the delivery of oxygen within the circulatory system is certainly advantageous. Hemocyanin has been identified in Onychophora and thus was present in the last common ancestor of Arthropoda (Burmester 2002; Kusche et al. 2002). It is extremely unlikely that hemocyanin was reinvented in Remipedia. It must be assumed that this respiratory protein was present in the last common ancestor of Malacostraca and Remipedia and was the principal oxygen carrier of stemline crustaceans. Thus, the occurrence of hemocyanin is a plesiomorphic character of Crustacea. As mentioned above, most non-malacostracan crustaceans use hemoglobin not hemocyanin for oxygen transport (Mangum 1985). However, this hemoglobin is a secondary invention (apomorphic character), which most likely derived from an intracellular globin of unknown function. It is currently uncertain whether hemoglobin emerged only once in the Crustacea, which would provide evidence that at least part of the “Entomostraca” sensu Walossek (1999), that is, Branchiopoda and Maxillopoda, may represent a monophylum.

Hexapod Affinities of Remipede Hemocyanins

Surprisingly, Blast searches and pairwise sequence comparisons revealed that the hemocyanin subunits of *S. tulumensis* display the highest sequence identity with insect hemocyanin subunits. For example, StuHc1 and StuHc3 share 60.4% and 53.4%, respectively, of the amino acids with the hemocyanin 1 of the firebrat *T. domestica*, whereas StuHc2 and the hemocyanin 2 of the cockroach *Blaptica dubia* are 57.1% identical. At least 10% lower identity scores were observed when the *S. tulumensis* hemocyanins were compared with those of malacostracan crustaceans or other arthropod hemocyanins. This evidence is a first rough indicator of a close relationship of hexapod and remipede hemocyanins.

The amino acid sequences of Remipedia hemocyanins were included in an alignment of arthropod hemocyanins and related proteins (Burmester and Scheller 1996; Burmester 2002; Hagner-Holler et al. 2004; Pick et al. 2009) (supplementary fig. 2, Supplementary Material online). Phylogenetic analyses employing ML and Bayesian methods resulted in essentially identical trees (fig. 3; supplementary figs. 3 and 4, Supplementary Material online).

Speleonectes tulumensis hemocyanins and hexapod proteins form a monophyletic clade that excludes other crustacean (malacostracan) sequences (ML bootstrap support: 48%; Bayesian posterior probability: 0.96). The other crustacean hemocyanins and pseudohemocyanins were monophyletic (100%; 1.0), but in none of the trees did the hemocyanins of the Remipedia join this clade. The relative position of the myriapod hemocyanins remains uncertain and received poor support. These proteins were either associated with the chelicerate hemocyanins (thereby supporting the “Paradoxopoda” hypothesis; Mallatt et al. 2004) or are in sister-group position to the crustacean plus hexapod proteins (supporting the traditional “Mandibulata”). In none of the trees did the myriapod hemocyanins join hexapod hemocyanins and hexamerins.

Speleonectes tulumensis hemocyanin subunits 1 and 3 (StuHc1 and StuHc3) form a common branch. This is reasonably well supported in Bayesian analyses (0.95) but only in 46% of the ML bootstrapped trees (fig. 3). A common clade of StuHc1 and StuHc3 and hexapod hemocyanin type 1 subunits plus hexamerins was recovered with 1.00 posterior probability in the Bayesian tree and supported by an ML bootstrap value of only 54%. The association of StuHc2 with the hexapod hemocyanin subunits 2 receives high support throughout (97%; 1.00) and is further corroborated by a unique and conserved insertion of nine amino acids in beta-sheet 3A (amino acid position 416–420 in StuHc2; fig. 1). Such insertion is not present in any other hemocyanin.

Tree topologies were further evaluated by hypothesis testing (table 1). The multiple ratio tests recovered the tree topology presented in figure 3 as best result and significantly reject StuHc1, StuHc2, and StuHc3 from the clade of other crustacean (malacostracan) hemocyanins. Hence, there is substantial molecular phylogenetic and structural evidence that the remipede hemocyanin subunits are orthologs of hexapod subunits. Thus, the lineage leading to the remipede and hexapod hemocyanins split into two distinct subunit types before these taxa diverged.

Are Remipedia and Hexapoda Sister Groups?

The origin of Hexapoda is notoriously disputed. Based on a large number of morphological characters, it has long been assumed that Hexapoda evolved from a myriapod-like ancestor and that these taxa form the subphylum “Tracheata” or “Atelocerata” (e.g., Brusca and Brusca 2003). However, this view has been challenged by molecular phylogenetic approaches that have provided evidence that Hexapoda are somehow, in fact, allied with Crustacea (Friedrich and Tautz 1995; Boore et al. 1998; Hwang et al. 2001; Kusche and Burmester 2001; Mallatt et al. 2004). Such topology was recovered in our analyses as well (fig. 3). A relationship of Hexapoda and Crustacea also received support in some comparative morphological and developmental biology studies (Giribet and Ribera 2000; Richter 2002), and thus Hexapoda and Crustacea have been joined in a common taxon named either “Tetraconata” (Richter 2002) or “Pancrustacea” (Zrzavý and Štys 1997).

In many analyses, Crustacea form a paraphyletic assemblage with respect to Hexapoda, but it has remained




FIG. 3.—Simplified Bayesian phylogenetic tree of arthropod hemocyanins and hexamerins. The numbers at the nodes show ML nonparametric bootstrap support (above) and Bayesian posterior probabilities (below). The complete trees are shown in supplementary figures 3 and 4, Supplementary Material online.

uncertain which crustacean taxon is the sister group of the Hexapoda. In most studies, only two crustacean classes were considered: Malacostraca and Branchiopoda. Analyses employing mitochondrial genomes suggest a close relationship of Malacostraca and Hexapoda with exclusion of Branchiopoda (Garcia-Machado et al. 1999; Wilson et al. 2000; Hwang et al. 2001). Such topology is also supported by some morphological evidence such as brain structure (Harzsch 2002) and by comparative studies of embryonic

development (Averof and Akam 1995). On the other hand, a clade consisting of Branchiopoda and Hexapoda tends to be supported by rRNA genes and multigene analyses (Regier et al. 2005, 2008; Mallatt and Giribet 2006; Roeding et al. 2007; Dunn et al. 2008).

Here, we provide molecular evidence that actually the Remipedia may be the closest living relatives of the Hexapoda. This relationship is suggested in our study 1) by the presence of both hexapod-type hemocyanin subunits in

Table 1
—Statistical Hypothesis Tests

Tree ^a	Log <i>L</i>	Difference	SE	SH ^b	1sKH ^b	2sKH ^b	ELW ^b
1	-74,424.60	0.00	Best	1.0000 +	1.0000 +	Best	0.9983 +
2	-74,509.58	84.98	21.5520	0.0000 -	0.0000 -	—	0.0000 -
3	-74,469.85	45.25	12.0780	0.0410 -	0.0020 -	—	0.0017 -
4	-74,460.23	35.63	12.6676	0.0120 -	0.0000 -	—	0.0000 -

^a Tree #1: Optimal tree as presented in figure 3: StuHc1 and 3 associated with type 1 hexapod hemocyanins and StuHc2 associated with type 2 hexapod hemocyanins. Tree #2: StuHc1, 2, and 3 in sister-group position to the other crustacean (malacostracan) hemocyanins. Tree #3: StuHc1 and 3 in sister-group position to the other crustacean (malacostracan) hemocyanins. Tree #4: StuHc2 in sister-group position to the other crustacean (malacostracan) hemocyanins.

^b Plus signs (+) denote the confidence sets, minus signs (-) denote rejection of a tree. All tests used 5% significance level.

Remipedia (but absence of any other subunit type), 2) by a unique sequence motif insertion that is shared by StuHc2 and hexapod-type 2 subunits, and 3) by the orthology of remipede and hexapod hemocyanin subunits (fig. 3), which is also supported by statistical tests of alternative tree topologies. A sister-group relationship of Remipedia and Hexapoda was tentatively retrieved in another molecular tree using elongation factor 2 sequences (Regier and Shultz 2001) and was also suggested by Telford et al. (2008) on the basis of unpublished material. In a large-scale phylogenomic study by Regier et al. (2008), a clade consisting of Remipedia and Hexapoda was consistently recovered, which in that study also included the Cephalocarida (which are not considered here).

Interestingly, a recent morphological study supports a close relationship of Remipedia and Hexapoda: Based on the structure of the arthropod brain, Fanenbruck and coworkers (Fanenbruck et al. 2004) proposed a common clade of Malacostraca, Remipedia, and Hexapoda (although they did not include Myriapoda in their analyses) with the exclusion of Branchiopoda and Maxillopoda. It was noted that the arrangement of nerves, axonal tracts, neuropil compartments and cell clusters in the brains of these taxa are similar but distinct from that of any other crustacean class. This hypothesis is now reinforced by the fact that all three taxa employ hemocyanin as respiratory protein, whereas other crustacean taxa use hemoglobins (Mangum 1985; Markl and Decker 1992; Burmester 2002). Therefore, Pancrustacea may be divided into two clades consisting of the possibly monophyletic Entomostraca (which have lost hemocyanins) on one hand (Walossek 1999), and Malacostraca, Remipedia, and Hexapoda (taxon N.N.) on the other (Fanenbruck et al. 2004). However, it should be noted that we cannot reject the hypothesis of multiple independent losses of hemocyanin and independent evolution of hemoglobin in various “entomostracan” lineages. If this is the case and the cited morphological evidence is ignored, the minimal topology recovered in our analyses is (Malacostraca, (Remipedia, Hexapoda)). Then an entomostracan, possibly branchiopod, ancestor of Hexapoda cannot be excluded (Regier et al. 2005; Mallatt and Giribet 2006; Roeding et al. 2007; Dunn et al. 2008).

Conclusions: Remipedia-Like Ancestor of Hexapoda?

Although recent hypotheses focused on a branchiopod-like crustacean as possible ancestor of hexapods (Glenner et al. 2006), our findings suggest that alternatively a remipede ancestor could be proposed. Extant and fossil Remipedia may provide evidence for the *Bauplan* of the hexapod stem lineage. The oldest known hexapod fossil is the collembolan *Rhyniella praecursor* from the Devonian Rhynie *Lagerstätte* in Scotland. This species harbors various derived features, thus offering little information on early morphology of Hexapoda. Interestingly, Haas et al. (2003) noted that *Tesnusocaris goldichi*, a remipedian fossil from the Carboniferous (Emerson and Schram 1991) may be a stem-lineage hexapod. This proposal was based on the unbranched structure of the first antenna and the long and filiform caudal appendages. It may be further speculated that

the enigmatic fossil *Devonohexapodus bocksbergensis* (Haas et al. 2003), which was described as a marine hexapod from the lower Devonian, may represent a transitional form between Remipedia and Hexapoda. Hexapod-like structures include leg-like palps of maxillae, absence of a second pair of antennae, and three pairs of longer uniramous thoracopods with six podomeres, but the homonomous trunk with its 38 trunk segments and the “abdominal” leglets actually provide an overall Remipedia-like appearance. In summary, it is possible that both fossil and living Remipedia occupy a key position for unraveling the evolution of Pancrustacea and thus for the understanding of morphological and functional innovations eventually resulting in the emergence of Hexapoda. However, additional molecular and morphological studies are required to unravel the position of Remipedia and to stably resolve their relationship to Hexapoda.

Supplementary Material

Supplementary table 1 and supplementary figures 1–4 are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

Acknowledgments

We wish to thank Christian Pick for providing arthropod hemocyanin sequences and advice. We are grateful to Tom Iliffe, Brett Gonzalez, and Jesse Henson for their help with collecting the *S. tulumensis* specimens. This work is supported by the Deutsche Forschungsgemeinschaft (Bu956/8, Bu956/9, Mi649/6, and Wa530/34) within the priority project 1174 “Deep Metazoan Phylogeny.”

Literature Cited

- Abascal F, Zardoya R, Posada D. 2005. ProfTest: selection of best-fit models of protein evolution. *Bioinformatics*. 21:2104–2105.
- Averof M, Akam M. 1995. Hox genes and the diversification of insect and crustacean body plans. *Nature*. 376:420–423.
- Boore JL, Lavrov DV, Brown WM. 1998. Gene translocation links insects and crustaceans. *Nature*. 392:667–668.
- Brusca RC, Brusca GJ. 2003. *Invertebrates*. Sunderland (MA): Sinauer Associates.
- Burmester T. 2001. Molecular evolution of the arthropod hemocyanin superfamily. *Mol Biol Evol*. 18:184–195.
- Burmester T. 2002. Origin and evolution of arthropod hemocyanins and related proteins. *J Comp Physiol [B]*. 172:95–107.
- Burmester T, Scheller K. 1996. Common origin of arthropod tyrosinase, arthropod hemocyanin, insect hexamerin, and dipteran arylphorin receptor. *J Mol Evol*. 42:713–728.
- Carapelli A, Lio P, Nardi F, van der Wath E, Frati F. 2007. Phylogenetic analysis of mitochondrial protein coding genes confirms the reciprocal paraphyly of Hexapoda and Crustacea. *BMC Evol Biol*. 7(Suppl 2):S8.
- Dunn CW, Hejnol A, Matus DQ, et al. (17 co-authors). 2008. Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature*. 452:745–749.
- Emerson MJ, Schram FR. 1991. Remipedia, part 2: paleontology. *Proc San Diego Soc Nat Hist*. 7:1–52.

- Fanenbruck M, Harzsch S. 2005. A brain atlas of *Godzilliognomus frondosus* Yager, 1989 (Remipedia, Godzilliidae) and comparison with the brain of *Speleonectes tulumensis* Yager, 1987 (Remipedia, Speleonectidae): implications for arthropod relationships. *Arthropod Struc Dev.* 34:343–378.
- Fanenbruck M, Harzsch S, Wagele JW. 2004. The brain of the Remipedia (Crustacea) and an alternative hypothesis on their phylogenetic relationships. *Proc Natl Acad Sci USA.* 101:3868–3873.
- Friedrich M, Tautz D. 1995. Ribosomal DNA phylogeny of the major extant arthropod classes and the evolution of myriapods. *Nature.* 376:165–167.
- Garcia-Machado E, Pempera M, Dennebouy N, Oliva-Suarez M, Mounolou JC, Monnerot M. 1999. Mitochondrial genes collectively suggest the paraphyly of Crustacea with respect to Insecta. *J Mol Evol.* 49:142–149.
- Gaykema WPI, Hol WGJ, Vereijken JM, Soeter NM, Bak HJ, Beintema JJ. 1984. 3.2 Å structure of the copper-containing, oxygen-carrying protein *Panulirus interruptus* haemocyanin. *Nature.* 309:23–29.
- Giribet G, Edgecombe GD, Wheeler WC. 2001. Arthropod phylogeny based on eight molecular loci and morphology. *Nature.* 413:157–161.
- Giribet G, Ribera C. 2000. A review of arthropod phylogeny: new data based on ribosomal DNA sequences and direct character optimization. *Cladistics.* 16:204–231.
- Glennier H, Thomsen PF, Hebsgaard MB, Sorensen MV, Willerslev E. 2006. The origin of insects. *Science.* 314:1883–1884.
- Goldman N, Anderson JP, Rodrigo AG. 2000. Likelihood-based tests of topologies in phylogenetics. *Syst Biol.* 49:652–670.
- Haas F, Waloszek D, Hartenberger R. 2003. *Devonohexapodus bocksbergensis*, a new marine hexapod from the Lower Devonian Hunsrück Slates, and the origin of Atelocerata and Hexapoda. *Org Divers Evol.* 3:39–54.
- Hagner-Holler S, Schoen A, Erker W, Marden JH, Rupprecht R, Decker H, Burmester T. 2004. A respiratory hemocyanin from an insect. *Proc Natl Acad Sci USA.* 101:871–874.
- Harzsch S. 2002. The phylogenetic significance of crustacean optic neuropils and chiasmata: a re-examination. *J Comp Neurol.* 453:10–21.
- Hassanin A. 2006. Phylogeny of Arthropoda inferred from mitochondrial sequences: strategies for limiting the misleading effects of multiple changes in pattern and rates of substitution. *Mol Phylogenet Evol.* 38:100–116.
- Hazes B, Magnus KA, Bonaventura C, Bonaventura J, Dauter Z, Kalk KH, Hol WG. 1993. Crystal structure of deoxygenated *Limulus polyphemus* subunit II hemocyanin at 2.18 Å resolution: clues for a mechanism for allosteric regulation. *Prot Sci.* 2:597–619.
- Herberts C, de Frescheville J. 1981. Occurrence of hemocyanin in the rhyzocephalan crustacea *Sacculina carcini* Thompson. *Comp Biochem Physiol B.* 70:657–659.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: bayesian inference of phylogenetic trees. *Bioinformatics.* 17:754–755.
- Hwang UW, Friedrich M, Tautz D, Park CJ, Kim W. 2001. Mitochondrial protein phylogeny joins myriapods with chelicerates. *Nature.* 413:154–157.
- Ito T. 1989. Origin of the basis in copepod limbs, with reference to remipedian and cephalocarid limbs. *J Crust Biol.* 9:85–103.
- Jaenicke E, Decker H, Gebauer W, Markl J, Burmester T. 1999. Identification, structure, and properties of hemocyanins from diplopod Myriapoda. *J Biol Chem.* 274:29071–29074.
- Katoh K, Kuma K, Toh H, Miyata T. 2005. MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res.* 33:511–518.
- Kishino H, Hasegawa M. 1989. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequence data, and the branching order in hominoidea. *J Mol Evol.* 29:170–179.
- Koenemann S, Olesen J, Alwes F, Iliffe TM, Hoenemann M, Ungerer P, Wolff C, Scholtz G. 2009. The post-embryonic development of Remipedia (Crustacea)—additional results and new insights. *Dev Genes Evol.* 219:131–145.
- Koenemann S, Schram FR, Bloechl A, Iliffe TM, Hoenemann M, Held C. 2007. Post-embryonic development of remipede crustaceans. *Evol Dev.* 9:117–121.
- Koenemann S, Schram FR, Hoenemann M, Iliffe TM. 2007. Phylogenetic analysis of Remipedia (Crustacea). *Org Divers Evol.* 7:33–51.
- Koenemann S, Schram FR, Iliffe TM. 2006. Trunk segmentation in Remipedia. *Crustaceana.* 79:607–631.
- Koenemann S, Schram FR, Iliffe TM, Hinderstein LM, Bloechl A. 2007. The behavior of Remipedia (Crustacea), with supporting field observations. *J Crust Biol.* 27:534–542.
- Kusche K, Burmester T. 2001. Diplopod hemocyanin sequence and the phylogenetic position of the Myriapoda. *Mol Biol Evol.* 18:1566–1573.
- Kusche K, Hembach A, Hagner-Holler S, Gebauer W, Burmester T. 2003. Complete subunit sequences, structure and evolution of the 6 × 6-mer hemocyanin from the common house centipede, *Scutigera coleoptrata*. *Eur J Biochem.* 270:2860–2868.
- Kusche K, Ruhberg H, Burmester T. 2002. A hemocyanin from the Onychophora and the emergence of respiratory proteins. *Proc Natl Acad Sci USA.* 99:10545–10548.
- Lavrov DV, Brown WM, Boore JL. 2004. Phylogenetic position of the Pentastomida and (pan)crustacean relationships. *Proc Biol Sci.* 271:537–544.
- Lim JT, Hwang UW. 2006. The complete mitochondrial genome of *Pollicipes mitella* (Crustacea, Maxillopoda, Cirripedia): non-monophyly of Maxillopoda and Crustacea. *Mol Cells.* 22:314–322.
- Mallatt JM, Garey JR, Shultz JW. 2004. Ecdysozoan phylogeny and Bayesian inference: first use of nearly complete 28S and 18S rRNA gene sequences to classify the arthropods and their kin. *Mol Phylogenet Evol.* 31:178–191.
- Mallatt JM, Giribet G. 2006. Further use of nearly complete 28S and 18S rRNA genes to classify Ecdysozoa: 37 more arthropods and a kinorhynch. *Mol Phylogenet Evol.* 40:772–794.
- Mangum CP. 1985. Oxygen transport in invertebrates. *Am J Physiol.* 248:505–514.
- Markl J, Decker H. 1992. Molecular structure of the arthropod hemocyanins. *Adv Comp Environ Physiol.* 13:325–376.
- Nicholas KB, Nicholas HBJ, Deerfield DWI. 1997. GeneDoc: analysis and visualization of genetic variation. *EMBnet NEWS* 4:14.
- Nielsen H, Engelbrecht J, Brunak S, von Heijne G. 1997. A neural network method for identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Int J Neural Syst.* 8:581–599.
- Pick C, Hagner-Holler S, Burmester T. 2008. Molecular characterization of hemocyanin and hexamerin from the firebrat *Thermobia domestica* (Zygentoma). *Insect Biochem Mol Biol.* 38:977–983.
- Pick C, Schneuer M, Burmester T. 2009. The occurrence of hemocyanin in Hexapoda. *FEBS J.* 276:1930–1941.
- Pohlmann JW, Iliffe TM, Cifuentes LA. 1997. A stable isotope study of organic cycling and the ecology of an anchialine cave ecosystem. *Mar Ecol Prog Ser.* 155:17–27.
- Regier JC, Shultz JW. 2001. Elongation factor-2: a useful gene for arthropod phylogenetics. *Mol Phylogenet Evol.* 20:136–148.

- Regier JC, Shultz JW, Ganley AR. (10 co-authors). 2008. Resolving arthropod phylogeny: exploring phylogenetic signal within 41 kb of protein-coding nuclear gene sequence. *Syst Biol.* 57:920–938.
- Regier JC, Shultz JW, Kambic RE, et al. 2005. Pancrustacean phylogeny: hexapods are terrestrial crustaceans and maxillopods are not monophyletic. *Proc Biol Sci.* 272:395–401.
- Richter S. 2002. The Tetraconata concept: hexapod-crustacean relationships and the phylogeny of Crustacea. *Organ Divers Evol.* 2:217–237.
- Roeding F, Hagner-Holler S, Ruhberg H, Ebersberger I, von Haeseler A, Kube M, Reinhardt R, Burmester T. 2007. EST sequencing of Onychophora and phylogenomic analysis of Metazoa. *Mol Phylogenet Evol.* 45:942–951.
- Schmidt HA, Strimmer K, Vingron M, von Haeseler A. 2002. TREE-PUZZLE: maximum likelihood phylogenetic analysis using quartets and parallel computing. *Bioinformatics.* 18:502–504.
- Schram FR. 1986. Crustacea. New York: Oxford University Press.
- Schram FR, Lewis CA. 1989. Functional morphology of feeding in Nectiopoda. In: Felgenhauer BE, Watling L, Thistle AB, editors. Crustacean issues 6: functional morphology of feeding and grooming in Crustacea. Rotterdam (Netherlands): A.A. Balkema Press. p. 115–122.
- Shimodaira H, Hasegawa M. 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol Biol Evol.* 16:1114–1116.
- Spears T, Abele LG. 1997. Crustacean phylogeny inferred from 18S rDNA. In: Fortey RA, Thomas RH, editors. Arthropod relationships. London: Chapman and Hall. p. 169–187.
- Strimmer K, Rambaut A. 2002. Inferring confidence sets of possibly misspecified gene trees. *Proc Biol Sci.* 269:137–142.
- Telford MJ, Bourlat SJ, Economou A, Papillon D, Rota-Stabelli O. 2008. The evolution of the Ecdysozoa. *Philos Trans R Soc Lond B Biol Sci.* 363:1529–1537.
- van der Ham JL, Felgenhauer BE. 2007. The functional morphology of the putative injecting apparatus of *Speleonectes tanumekes* (Remipedia). *J Crust Biol.* 27:1–9.
- van der Ham JL, Felgenhauer BE. 2008. Ultrastructure and functional morphology of glandular setae and distal claws of cephalic appendages of *Speleonectes tanumekes* (Crustacea: remipedia). *Arthropod Struct Dev.* 37:235–247.
- van Holde KE, Miller KI. 1995. Hemocyanins. *Adv Prot Chem.* 47:1–81.
- von Reumont BM, Meusemann K, Szucsich NU, et al. (13 co-authors). 2009. Can comprehensive background knowledge be incorporated into substitution models to improve phylogenetic analyses? A case study on major arthropod relationships. *BMC Evol Biol.* 9:119.
- Walossek D. 1999. On the Cambrian diversity of Crustacea. In: Schram FR, von Vaupel Klein JC, editors. Crustaceans and the biodiversity crisis, Proceedings of the 4th International Crustacean Congress. Leiden (Netherlands): Brill. p. 3–27.
- Whelan S, Goldman N. 2001. A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach. *Mol Biol Evol.* 18:691–699.
- Wills M. 1997. A phylogeny of recent and fossil Crustacea derived from morphological characters. In: Fortey RA, Thomas RH, editors. Arthropod relationships. London: Chapman & Hall. p. 189–209.
- Wilson K, Cahill V, Ballment E, Benzie J. 2000. The complete sequence of the mitochondrial genome of the crustacean *Penaeus monodon*: are malacostracan crustaceans more closely related to insects than to branchiopods? *Mol Biol Evol.* 17:863–874.
- Yager J. 1981. Remipedia, a new class of Crustacea from a marine cave in the Bahamas. *J Crust Biol.* 1:328–333.
- Yager J. 1991. The Remipedia (Crustacea): recent investigations of their biology and phylogeny. *Verh Dtsch Zool Ges.* 84: 261–269.
- Zrzavý J, Štys P. 1997. The basic body plan of arthropods: insights from evolutionary morphology and developmental biology. *J Evol Biol.* 10:353–367.

Billie Swalla, Associate Editor

Accepted August 15, 2009