

## Original article

# Ticks (Acari: Ixodidae) of three Timor-Leste reptiles: first country record of *Amblyomma helvolum*, with new interactions and an updated list of host species

Fabrício H. Oda <sup>a,\*</sup>, Thiago F. Martins <sup>b,c</sup>, Marcelo B. Labruna <sup>b</sup>, Mark O’Shea <sup>d</sup>, Hinrich Kaiser <sup>e,f</sup><sup>a</sup> Departamento de Química Biológica, Laboratório de Zoologia, Universidade Regional do Cariri, Campus Pimenta, Crato, Ceará, Brazil<sup>b</sup> Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, Brazil<sup>c</sup> Departamento de Laboratórios Especializados, Superintendência de Controle de Endemias, Secretaria de Estado da Saúde de São Paulo, São Paulo, Brazil<sup>d</sup> Faculty of Science and Engineering, University of Wolverhampton, Wulfruna Street, Wolverhampton WV1 1LY, United Kingdom<sup>e</sup> Sektion Herpetologie, Zentrum für Taxonomie und Morphologie, Museum Koenig, Leibniz-Institut zur Analyse des Biodiversitätswandels, Adenauerallee 160, 53113 Bonn, Germany<sup>f</sup> Department of Biology, Victor Valley College, 18422 Bear Valley Road, Victorville, California 92395, USA

## ARTICLE INFO

## ABSTRACT

## Keywords:

*Amblyomma helvolum**Carlia**Sphenomorphus**Coelognathus subradiatus*

Lesser Sunda Archipelago

Outer Banda Arc

We report on the presence of the ixodid tick *Amblyomma helvolum* on three species of reptiles from Timor-Leste. Among a total of 21 host specimens (lizards: 18 four-fingered skinks, *Carlia* sp. ‘Meleotegi’ and two forest skinks, *Sphenomorphus* sp. ‘Meleotegi’; snake: one *Coelognathus subradiatus*) four were parasitized by ticks. Whereas nymphs were associated with the lizards, an adult male was retrieved from the snake. This report is the first of *A. helvolum* for Timor-Leste, the first for the skink genera *Carlia* and *Sphenomorphus*, and the first for *C. subradiatus*. We present a comprehensive table with updated taxonomy of known associations between *A. helvolum* and other vertebrates.

## 1. Introduction

Timor-Leste, Asia’s newest country, is positioned at the south-central extreme of the Indonesian Archipelago, where it occupies the eastern half of Timor Island (Fig. 1). Timor-Leste’s territory is composed of four landmasses, including the eastern half of the island, where 12 of the country’s 13 municipalities lie, the Oecusse exclave surrounded by Indonesian West Timor, the coralogenic Jaco Island at its eastern end, all within the Outer Banda Arc, and Ataúro Island, the only portion of Timor-Leste that lies in the Inner Banda Arc. The country’s position at the southeastern extreme of the biogeographic province known as Wallacea places it at the crossroads of four interesting landmasses: Southeast Asia and the islands of the Sunda Shelf to the west, Australia to the south, Sahul Shelf islands and Melanesia to the east, and Sulawesi to the north, all of which appear to have some influence on the terrestrial fauna, including reptiles (Monk et al., 1997; O’Shea et al., 2015). An evaluation of parasite groups is therefore an important piece to the biogeographic puzzle that is the fauna of Timor.

During ten herpetofaunal surveys (Kaiser et al., 2011; O’Shea et al.,


2012; Sanchez et al., 2012; Kaiser et al., 2013) HK and MOS and the members of the Tropical Research Initiative collected a variety of reptile species, some of which harbored ectoparasites, including ticks. In the following paragraphs, we detail the ixodid fauna recovered, report the first country record of *Amblyomma helvolum* Koch, 1844, and add some new host records for this species.

## 2. Materials and methods

## 2.1. Host survey

Lizard hosts were collected at two sites in close proximity to each other (Fig. 1) and covering an elevational range of 1150–1250 m. The first site is the Sta. Bakhita Mission in Eraúlo, Ermera Municipality (8.781°S, 125.4438°E), where two species of four-fingered skinks (genus *Carlia*) are sympatric (16S rRNA, CO1 mtDNA; Kaiser et al., unpubl. data). Two of the host specimens (USNM 579745, 581104), adult males collected on 5 February 2012 and 29 June 2013, respectively, are part of a north coast Timor population of *Carlia* that reaches its highest elevations in central

\* Corresponding author at: Departamento de Química Biológica, Laboratório de Zoologia, Universidade Regional do Cariri, Campus Pimenta, Crato, Ceará, Brazil.  
E-mail address: [fabricio\\_oda@hotmail.com](mailto:fabricio_oda@hotmail.com) (F.H. Oda).




**Fig. 1.** Map of Timor-Leste with labelled localities where tick specimens were collected from three reptile species. The red dot indicates the collection site of four-fingered skinks (*Carlia* sp. "Meleotegi") at the Santa Bakhita Mission, the yellow dot is the Meleotegi River collection site of *Carlia* sp. "Meleotegi" and forest skinks (*Sphenomorphus* sp. "Meleotegi"), and the pink dot is the locality where a freshly killed Lesser Sunda Racer (*Coelognathus subradiatus*) was found dead on the road. The white rectangle in the inset globe shows the position of Map A. Map A shows the Lesser Sunda Islands with two island arcs, Inner and Outer Banda Arc. The scale bar in the main map represents a distance of 100 km. Beneath the maps, habitats of the collected reptiles are illustrated with frame colors corresponding to the colors of the locality dots. Photos by Mark O'Shea.

Ermera Municipality. This is an undescribed species, hereafter referred to as *Carlia* sp. 'Meleotegi' (Fig. 2A). The first specimen was collected in the grounds of the mission among garden plants, whereas the second was caught along the Meleotegi River ( $8.7806^{\circ}$ S,  $125.4543^{\circ}$ E), a rocky riverbed lined on both banks by agricultural plots, including coffee plantations. The second host, a member of an undescribed species of forest skink, genus *Sphenomorphus* (USNM 579765; Fig. 3A), was collected in the coffee forest on the western bank of the Meleotegi River on 5 February 2012. This specimen belongs to a high-elevation population apparently endemic to Ermera Municipality. It is not conspecific with *S. melanopogon*, a south coast low-elevation species (Shea, 2012; Justin Bernstein, pers. comm.), and this population is referred to hereafter as *Sphenomorphus* sp. 'Meleotegi.' The snake host (USNM 580544; Fig. 4A) is a freshly killed specimen of the Lesser Sunda Racer (*Coelognathus subradiatus*), found dead on a major north–south road near Lacubar, Manatuto Municipality ( $8.7765^{\circ}$ S,  $125.9653^{\circ}$ E, elevation 870 m; Fig. 1) on 2 July 2012. Lizards were euthanized by injection with 5% procaine (Livezey, 1958) following standard animal care protocols (e.g., ASIH, 2004), fixed in 10% formalin, and stored in 70% ethanol. Animals were collected under the auspices of the Smithsonian Institution, Washington, D.C., U.S.A. with export permission from the Departamento Areas Protegidas e Parque Nacional, Direcção Nacional de Floresta, Secretario do Estado de Floresta no Konservasaun Natureza, Ministerio Agricultura e Pescas, Timor-Leste (permit nos. 26/DAPPN-DNF-MAP/I/2012, 22/DAPPN-DNF-MAP/VI/2012, 29/DAPPN-DNF-MAP/VI/2013) and


import permission from the United States Fish and Wildlife Service (Forms 3-177, clearance nos. 2012SF1125818, 2012HN1197038, 2013SF1361984).

## 2.2. Photography


The live skinks and snake were photographed on sets built in a portable 19 cm x 90 cm x 90 cm Lastolite Cubelite photography dome using Canon EOS 7D camera bodies in combination with either a Canon EF-S 60 mm f2.8 USM macro lens or a Canon EF 100 mm f2.8L IS USM macro lens. Illumination was provided by Canon Macro Twin Lite MT-24EX flash units. The ticks were photographed in situ and ex situ using the same camera body with a Canon MP-E 64 mm f2.8 1-5x macro lens and a Canon Macro Twin Lite MT-24EX flash unit. This lens was pushed to 10x life size by the addition of a Canon EF 2x Extender. Figures were prepared on a Mac Pro computer running OS 10.12 Sierra and Adobe Photoshop CC 2014. Parasite images were cut out and placed on a black background using a Wacom Cintiq 13-inch HD Touch connected to an iMac Pro computer running OS 11.5 Big Sur and Adobe Photoshop 2021. The Timor-Leste map was adapted from a Shutterstock Schwanenblitz 3D rendering, the world globe was prepared using Mountain High relief maps (<http://www.mountainhighmap.com>).



**Fig. 2.** An undescribed species of four-fingered skink, genus *Carlia*, from Timor-Leste and associated ticks. (A) An adult male *Carlia* sp. 'Meleotegi' with bright coloration indicating breeding readiness. (B) Engorged nymph of *Amblyomma helvolum* in a near-axillary position. (C) Another engorged nymph, showing the tick's capitulum deeply embedded in between two ventral scales. (D) Two *A. helvolum* nymphs after detachment from their host in dorsal (right) and ventral (left) views. Photos by Mark O'Shea.



**Fig. 3.** An undescribed species of forest skink, genus *Sphenomorphus*, from Timor-Leste and an associated tick. (A) An adult *Sphenomorphus* sp. 'Meleotegi' of undetermined sex showing the characteristic lateral striping of the species. (B) Engorged nymph of *Amblyomma helvolum* in an axillary position. Photos by Mark O'Shea.



**Fig. 4.** (A) A dead *Coelognathus subradiatus* photographed *in situ* on one of the main north-south roads in Timor-Leste. It is noteworthy that this snake was not run over by a vehicle, it was instead noosed, beaten to death, and thrown into the road by a local resident. (B) An adult *C. subradiatus* (USNM 579240) from Timor-Leste in life. (C) Adult male *Amblyomma helvolum* photographed *in situ* while attached to the snake in (A). (D) Dorsal and ventral views in life of an adult male *A. helvolum* after detachment from the snake. The dorsal view (left) shows the dorsal colour and pore pattern on the strongly demarcated shield of this species. The ventral view of the same individual (right) shows anus, anal groove, genital aperture, and the laterally positioned spiracular plates. Photos by Mark O'Shea.

### 2.3. Tick identification

Each host specimen was examined for the presence of ectoparasites immediately after euthanasia. Ticks were removed from host bodies

with forceps and preserved in 70% ethanol. The collected ticks were sent to the Department of Preventive Veterinary Medicine and Animal Health (VPS) of the School of Veterinary Medicine and Animal Science (FMVZ) at the University of São Paulo (USP), São Paulo, Brazil. For taxonomic

**Table 1**

Lizard and snake hosts of *Amblyomma helvolum* from Timor-Leste in the Lesser Sunda Archipelago. Column abbreviations include the number of examined hosts ( $n_e$ ), the number of infested hosts ( $n_i$ ), the number of collected ticks ( $n_c$ ), and the site of collection (Sta. Bakhita Mission, BM; Meleotegi River, MR; and Laclubar, LB).

Host Species	$n_e$	$n_i$	<i>Amblyomma helvolum</i> Stage	$n_c$	Site
<b>Lizards</b>					
<i>Carlia</i> sp. 'Meleotegi'	16	2	Nymphs	3	BM, MR
<i>Sphenomorphus</i> sp. 'Meleotegi'	11	1	Nymph	1	MR
<b>Snakes</b>					
<i>Coelognathus subradiatus</i>	8	1	Adult male	1	LB
<b>Total</b>	<b>35</b>	<b>4</b>		<b>5</b>	

identification we used the reports of Anastos (1950), Robinson (1926), and Voltzit and Keirans (2002). Tick voucher specimens were deposited in the tick collection "Coleção Nacional de Carrapatos Danilo Gonçalves Saraiva" (CNC) of the FMVZ-USP, under accession numbers CNC 4205-4207.

#### 2.4. Bibliographic review

A bibliographic search was conducted using Google Scholar to update the list of Simmons et al. (2002) of *A. helvolum* hosts. The search terms used were: "*Amblyomma helvolum*", "amphibians", "reptiles", "lizards", "snakes", "turtles", "birds", and "mammals". Old references cited by publications recovered by the primary bibliographic search were located using Biodiversity Heritage Library (<https://www.biodiversitylibrary.org/>), BioStor-Lite (<http://biostor.org/>), and Internet Archives (<https://archive.org/>). Scientific names of hosts were updated to current nomenclature following Uetz et al. (2021) (<http://www.reptile-database.org/>) for lizards and snakes, and GBIF.org (2020) | Global Biodiversity Information Facility (<https://www.gbif.org/>) for birds and mammals.

### 3. Results

During the surveys that are the subject of this study, 16 specimens of *Carlia* sp. 'Meleotegi,' eleven of *Sphenomorphus* sp. 'Meleotegi,' and eight of *C. subradiatus* were collected, totaling 35 host specimens. Of these, four host specimens (two *Carlia*, one *Sphenomorphus*, one *Coelognathus*) had ticks. A total of five ticks were retrieved from the hosts. Based on morphological characteristics described in the literature, the ticks were identified as four nymphs (Fig. 2D) and one adult male (Fig. 4C, D) of *A. helvolum*. Nymphs were recorded from *Carlia* sp. 'Meleotegi,' and *Sphenomorphus* sp. 'Meleotegi,' and the adult male from *C. subradiatus* (Table 1).

The adult *A. helvolum* male was identified by the following characteristics: scutum with eyes, without marginal groove, ornate, with yellowish patches that include large punctations in the scapular, marginal, and central areas; Coxa I with two spurs, the external twice as long as the internal, Coxae II-III with one triangular spur; basis capitulum rectangular, with small rounded cornua. The nymphs were characterized as follows: scutum with eyes and few small punctations; Coxa I with two spurs, the external longer, the internal small and situated anteriorly, coxae II-III with one triangular spur; basis capitulum triangular, without cornua.

The position of ticks on the bodies of their lizard hosts varied slightly. In *Carlia* sp. 'Meleotegi,' nymphs were attached in an axillary position on either the forelimb or the hind limb (Fig. 2B), but also on the exposed surfaces of the belly (Fig. 2C). In *Sphenomorphus* sp. 'Meleotegi,' an engorged nymph was found in an axillary attachment position (Fig. 3B), whereas in *C. subradiatus* the adult male was attached on the dorsal surface (Fig. 4C).

Forty-four studies have reported parasitism records of *A. helvolum* in nine countries. Most of the records are from Indonesia (30), the Philippines (24), and Malaysia (23), followed by Thailand (16), India

(13), Singapore (4), China, Laos, and Vietnam (one record each). *Amblyomma helvolum* was reported on 41 reptile species (15 lizards, 27 snakes, and one turtle), one bird species, and eight mammal species. Amongst the reptiles, varanids and colubrids were the most affected, accounting for 60% and 63% of the parasitism records reported in lizards and snakes, respectively (Table 2).

### 4. Discussion

*Amblyomma helvolum* is native to the Australasian and Oriental biogeographic realms, where it has been reported from China, India, Indonesia, Laos, Malaysia, the Philippines, and Vietnam (Auffenberg, 1988; Kolonin, 1995; Petney and Keirans, 1995; Ghosh et al., 2007; Chen et al., 2010; Chao et al., 2013). Australia and Papua New Guinea were included in the geographic distribution of *A. helvolum* by Burridge (2011), but its presence there has not been confirmed with voucher specimens (Guglielmone and Robbins, 2018).

The records of *A. helvolum* on two snakes in West Bengal, India, by Ghosh et al. (2018) are erroneous because of morphological discrepancies between those specimens and *A. helvolum*. Ghosh et al. (2018) described and illustrated the ticks retrieved from the two snakes as without eyes and Coxa I with internal and external spurs almost equal in size (eyes present, Coxa I with external spur about twice as long as the internal spur in *A. helvolum*; Anastos, 1950; Robinson, 1926; Voltzit and Keirans, 2002). We therefore follow Guglielmone et al. (2015) and consider the records of *A. helvolum* by Ghosh et al. (2017, 2021) as unconfirmed, since it seems that the authors of those reports used the same characteristics as adopted by Ghosh et al. (2018), and they are most likely of a type of tick that is not conspecific with *A. helvolum*. The remaining records of *A. helvolum* from West Bengal (India) are all from captive hosts (Table 2), and the presence of natural populations of this tick species in West Bengal remains to be confirmed.

There are three records of the introduction of *A. helvolum* into the United States through imported snakes from Thailand and Indonesia (USDA, 1971; Clark and Doten, 1995; Simmons et al., 2002; Keirans and Durden, 2001; Burridge and Simmons, 2003), but there is no evidence that this tick has become established there (Guglielmone et al., 2014). The tick fauna of Timor-Leste was inventoried for the first time during a parasitological survey in 1973, but *A. helvolum* was not found (Dias, 1988). Our finding, therefore, is the first record of *A. helvolum* for Timor-Leste.

Records of *A. helvolum* parasitizing squamates include species of the lizard families Agamidae, Scincidae, and Varanidae, and species of the snake families Boidae, Colubridae, Elapidae, Natricidae, and Pythonidae. Among these, there are no reports to date of *A. helvolum* parasitizing species of the lizard genera *Carlia* and *Sphenomorphus*, nor on the snake *C. subradiatus*. Parasitism of *A. helvolum* on the scincid lizards *Carlia* sp. 'Meleotegi' and *Sphenomorphus* sp. 'Meleotegi', and the colubrid snake *C. subradiatus* are new interactions for this tick species. The herpetofauna of Timor-Leste comprises at least seven species of anurans, 42 species of lizards, 17 species of snakes, two species of turtles, and one species of crocodile (O'Shea et al., 2011, 2015). The three reptile species parasitized by *A. helvolum* correspond to an unexpectedly low frequency

**Table 2**

Checklist of the reptile hosts of *Amblyomma helvolum*. Columns include the host species, the name of the tick given in the source publication (when a current synonym of *A. helvolum* was used), the life stage reported (adult with unspecified sex, A; adult male, M; adult female, F; nymphs, N; larvae, L; no information, x), the country of record, and the source publication. Country abbreviations include China (CHN), Indonesia (IDN), India (IND), Laos (LAO), Malaysia (MYS), the Philippines (PHL), Singapore (SGP), Taiwan (TWN), Thailand (THA), Timor-Leste (TLS), and Vietnam (VNM). Species names with an asterisk (\*) are those species listed as hosts of *A. helvolum* by Simmons et al. (2002).

Host	Name	Stage	Location (Country)	Source
<b>LIZARDS</b>				
<b>Agamidae</b>				
* <i>Gonocephalus sophiae</i> (Gray, 1845)		A	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
* <i>Hydrosaurus pustulatus</i> (Eschscholtz, 1829)		F,M	Davao Province, Mindanao (PHL)	Kohls (1950)
<b>Scincidae</b>				
* <i>Carlia</i> sp. 'Meleotegi'		N	Ermera Municipality (TLS)	This study
* <i>Dasia grisea</i> (Gray, 1845)		N	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
* <i>Eutropis multifasciata</i> (Kuhl, 1820)		x	Simeulue Island, Sumatra (IDN)	Warburton (1926)
		F,M	Java (IDN)	Krijgsman and Ponto (1931)
		F,N	Culion Island, Palawan Province (PHL)	Kohls (1950)
		F,N	Jarak Island, Sembilan Islands, Perak (MYS)	Audy et al. (1960)
		F,N,L	Rumbia Island, Sembilan Islands, Perak (MYS)	Audy et al. (1960)
		N,L	Lalang Island, Sembilan Islands, Perak (MYS)	Audy et al. (1960)
		N	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
		x	Krau Wildlife Reserve, Pahang (MYS)	Kho et al. (2019)
		A,N	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
		N	Ermera Municipality (TLS)	This study
		A	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
* <i>Otosaurus cumingii</i> Gray, 1845		F	Songkhla Province (THA)	Sharif (1928)
* <i>Sphenomorphus</i> sp. 'Meleotegi'		A	unknown locality (THA)	Tanksul et al. (1983)
* <i>Tropidophorus grayi</i> Günther, 1861		F,M	Srisaket Province (THA)	Kaenkan et al. (2020)
<b>Varanidae</b>				
* <i>Varanus bengalensis</i> (Daudin, 1802)		F,M,N	unknown locality (MYS)	Audy et al. (1960)
		x	no locality provided	King and Green (1999)
		F,M	Komodo and Flores (IDN)	Schulze (1933)
		x	Komodo (IDN)	Pianka et al. (2004)
		x	Panay (PHL)	Pianka et al. (2004)
		M	Manila, Luzon (PHL)	Neumann (1910)
		A	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988) [as <i>V. salvator</i> ]
		x	Luzon (PHL)	Parrish (1971) [as <i>Varanus</i> sp.]
		M	Negros Occidental Province (PHL)	Keirans (1984) [as <i>Varanus</i> sp.]
		F,M,N	Camarines Sur Province, Luzon (PHL)	Auffenberg (1988)
		x	Luzon (PHL)	Pianka et al. (2004)
		F	Palawan Province (PHL)	Kohls (1950) [as <i>V. salvator</i> ]
		M,N	unknown locality (MYS)	Audy et al. (1960)
		M	unknown locality (SGP)	Robinson (1926)
		x	unknown locality (SGP)	Kohls (1957)
<i>Varanus salvator macromaculatus</i>	<i>Amblyomma furcosum</i>	F	Sumatra (IDN)	Neumann (1910)
Deraniyagala, 1944				
<i>Amblyomma quadrivaculatum</i>				
<i>A. decoratum</i>				
		F,M	Sumatra (IDN)	Schulze (1933)
		F,M	Sumatra (IDN)	Neumann (1911)
		F	Simeulue Island, Sumatra (IDN)	Warburton (1926)
		x	Tioman Island, Pahang (MYS)	Kwak (2020)
		F	Pichit Province (THA)	Doornbos et al. (2013)
		x	Sabah (MYS)	Guerrero Sanchez (2019)
		F,M	Narathiwat and Pattani Provinces (THA)	Kaenkan et al. (2020)
		M	Nicobar Islands (IND)	Warburton (1910) [as <i>V. salvator</i> ]
		M	Nicobar Islands (IND)	Robinson (1926) [as <i>V. salvator</i> ]
		F,M	Java (IDN)	Krijgsman and Ponto (1931)
		M	Java (IDN)	Schulze (1933)
		F,M	Rumbia Island, Perak (MYS)	Audy et al. (1960)
		F,M	Java (IDN)	Keirans (1984)
		M	Tanimbar Islands (IDN)	Hirst and Hirst (1910)
<i>Varanus</i> sp. 'Tanimbar'	<i>Amblyomma (Aponomma) tenimberense</i>			
<i>Varanus</i> sp.		M	Sakijang Bendera Island (SGP)	Robinson (1926), Keirans (1984)
		F,M	Java (IDN)	Anastos (1950)
		A	unknown locality (THA)	Tanskul et al. (1983)
		F,M	West Bengal (IND)	Sanyal and De (1992) <sup>c</sup>
		F	unknown locality (LAO)	Petney et al. (1995)
<b>SNAKES</b>				
<b>Colubridae</b>				
* <i>Boiga dendrophila</i> (Boie, 1827)	<i>A. quadrivaculatum</i>	M	Java (IDN)	Neumann (1899)
		F,M	Berhala Island, Sumatra (IDN)	Oudemans (1928)
	<i>A. quadrivaculatum</i>	M	Berhala Island, Sumatra (IDN)	Schulze (1933)
		F	Sabah (MYS)	Mao et al. (2019)

(continued on next page)

**Table 2 (continued)**

Host	Name	Stage	Location (Country)	Source
* <i>Coelognathus flavolineatus</i> (Schlegel, 1837)		F,M,N	unknown locality (MYS)	Audy et al. (1960)
* <i>Coelognathus radiatus</i> (Boie, 1827)		F	Songkhla Province (THA)	Sharif (1928)
		A	unknown locality (THA)	Tanskul et al. (1983)
<i>Coelognathus subradiatus</i> (Schlegel, 1837)		M	Manatuto Municipality (TLS)	This study
<i>Dendrelaphis pictus</i> (Gmelin, 1789)		F	Songkhla Province (THA)	Keirans (1984)
<i>Elaphe carinata</i> (Günther, 1864)		F	Kaohsiung City (TWN)	Chao et al. (2013)
<i>Gonyosoma oxycephalum</i> (Boie, 1827)		F	Java (IDN)	Robinson (1926) <sup>a</sup> , Keirans (1984)
<i>Lycodon capucinus</i> (Boie, 1827)		F	Cagayan Province, Luzon (PHL)	Tabug and Necesito (2021)
<i>Ptyas carinata</i> (Günther, 1858)		F,M	Mahasarakham and Kalasin Provinces (THA)	Kaenkan et al. (2020)
* <i>Ptyas dipsas</i> (Schlegel, 1837)		M	Sulawesi (IDN)	Lazell et al. (1991)
* <i>Ptyas fusca</i> (Günther, 1858)		F,N,L	unknown locality (MYS)	Audy et al. (1960)
* <i>Ptyas korros</i> (Schlegel, 1837)		F	Kuala Lumpur, Federal State (MYS)	Robinson (1926), Keirans (1984)
		F,M	Java (IDN)	Robinson (1926), Keirans (1984)
		F,M	Mindanao (PHL)	Nakatsuji (1937)
		F	Java (IDN)	Anastos (1950)
		x	unknown locality ("Southeast Asia")	Dias (1958)
		F	Nam Phong District, Khon Kaen Province (THA)	Sumrandee et al. (2014a, 2014b, 2015)
* <i>Ptyas mucosa</i> (Linnaeus, 1758)	<i>A. decoratum</i>	F,M	Sumatra (IDN)	Neumann (1911)
	<i>Amblyomma feuerborni</i>	F,M	Klakah, Java (IDN)	Schulze (1933)
		F,M,N	unknown locality (TWN)	Chao et al. (2013)
<b>Elapidae</b>				
* <i>Naja naja</i> (Linnaeus, 1758)		M	Selangor (MYS)	Kohls (1957)
		F	unknown locality (MYS)	Audy et al. (1960)
<i>Naja sumatrana</i> Müller, 1887		F,M	West Bengal (IND)	Sanyal and De (1992) <sup>c</sup>
		F	Sumatra (IDN)	Anastos (1950)
		M	Selangor (MYS)	Kohls (1957) [as <i>N. naja</i> ]
		x	Johor (MYS)	Kho et al. (2015)
		x	Babi Island, Riau Archipelago (IDN)	Warburton (1926)
<b>Natricidae</b>				
* <i>Ophiophagus hannah</i> (Cantor, 1836)		F,M	Srisaket Province (THA)	Kaenkan et al. (2020)
<i>Fowlea flavipunctatus</i> (Hallowell, 1860)		F	Phichit Province (THA)	Sumrandee et al. (2014a, 2014b)
<i>Fowlea piscator</i> (Schneider, 1799)		N	unknown locality (MYS)	Audy et al. (1960)
* <i>Rhabdophis flavigeeps</i> (Duméril et al., 1854)		F	unknown locality (MYS)	Audy et al. (1960)
* <i>Xenochrophis trianguligerus</i> (Boie, 1827)				
<b>Pythonidae</b>				
* <i>Malayopython reticulatus</i> (Schneider, 1801) <sup>b</sup>	<i>A. furcosum</i>	F	Java (IDN)	Neumann (1901)
	<i>A. quadrivaculatum</i>	F,M	Java (IDN)	Neumann (1901)
	<i>A. feuerborni</i>	M	Java (IDN)	Schulze (1933)
		x	unknown locality (PHL)	Nakatsuji (1937)
		F,M,N	Davao Province, Mindanao (PHL)	Kohls (1950)
		F,M,N,	unknown locality (MYS)	Audy et al. (1960)
		L		
		F	Đà Nẵng (VNM)	Hoogstraal et al. (1968)
		A	unknown locality (THA)	Tanskul et al. (1983)
<i>Python bivittatus</i> Kuhl, 1820		x	Lorong Halus, Singapore island (SGP)	Kim (2020)
		F	Nakhon Nayok Province (THA)	Sumrandee et al. (2014a, 2014b, 2015)
<i>Python</i> sp.		A	Selangor (MYS)	Kho et al. (2015) [as <i>P. molurus</i> ]
Pythonidae incertae sedis		x	Sarawak (MYS)	Robinson (1926), Keirans (1984)
		A	unknown locality (THA)	Tanskul et al. (1983)
<b>TURTLES</b>				
<b>Geoemydidae</b>				
* <i>Heosemys grandis</i> (Gray, 1860)	<i>A. decoratum</i>	M	West Bengal (IND)	Warburton (1910) <sup>c</sup> , Robinson (1926) <sup>c</sup>
		x	West Bengal (IND)	Sanyal and De (1992) <sup>c</sup>
<b>BIRDS</b>				
<b>Pittidae</b>				
<i>Erythropitta erythrogaster</i> (Temminck, 1823)		x	Luzon (PHL)	Parrish (1971)
<b>MAMMALS</b>				
<b>Bovidae</b>				
<i>Bubalus bubalis</i> (Linnaeus, 1758)		F	Sulawesi (IDN)	Durden et al. (2008)
<b>Cervidae</b>				
<i>Rusa marianna</i> (Desmarest, 1822)		x	Luzon (PHL)	Parrish (1971)
<b>Muridae</b>				
<i>Chrotomys whiteheadi</i> Thomas, 1895		x	Luzon (PHL)	Parrish (1971)
<i>Leopoldamys sabanus</i> (Thomas, 1887)		L	Tioman Island, Pahang (MYS)	Nadchatram et al. (1966) <sup>b</sup>
<i>Rattus everetti</i> Günther, 1879		x	Luzon (PHL)	Parrish (1971)
<i>Rattus rattus mindanensis</i> Mearns, 1905		x	Luzon (PHL)	Parrish (1971)
<i>Rattus tiomanicus</i> (Miller, 1900)		L	Tioman Island, Pahang (MYS)	Nadchatram et al. (1966) <sup>b</sup>
<b>Suidae</b>				
<i>Sus scrofa</i> Linnaeus, 1758		M	Sulawesi (IDN)	Durden et al. (2008)

<sup>a</sup> Listed as *Coluber onicephalus*.<sup>b</sup> Guglielmone et al. (2018) stated that the diagnoses of larvae of *A. helvolum* on Muridae require confirmation.<sup>c</sup> Host-parasite association in captive habitats.

(< 6%) of reptile species known to inhabit Timor, demonstrating that further studies are needed to better characterize the tick fauna and its incidence on amphibians and reptiles in this country.

Reptilian hosts of the tick *A. helvolum* were most recently reviewed by Simmons et al. (2002), who listed 26 species (13 lizards, 12 snakes, and one turtle), but not always employing the correct taxonomic allocation of host species. In that study, 21 reptile species were identified as additional records (Simmons et al., 2002). Our new host records and additional records increase the number of reptile species known as hosts of *A. helvolum* to 42 (Table 2). Water monitors (*Varanus salvator*) and reticulated pythons (*Malayopython reticulatus*) are the most common reptilian hosts of *A. helvolum* in Asia (Auffenberg, 1988; King and Green, 1999; Simmons et al., 2002). The records of *A. helvolum* from the literature for the scincid lizard *Eutropis multifasciata*, the genus *Varanus*, the colubrid snake *Lycodon capucinus*, and the python *Malayopython reticulatus* suggest that these species may also be hosts for *A. helvolum* in Timor-Leste, but their potential association there requires confirmation.

#### CRediT authorship contribution statement

**Fábrício H. Oda:** Conceptualization, Methodology, Writing – original draft, Writing – review & editing. **Thiago F. Martins:** Conceptualization, Methodology, Data curation, Writing – original draft, Writing – review & editing. **Marcelo B. Labruna:** Methodology, Data curation, Writing – original draft, Writing – review & editing. **Mark O'Shea:** Conceptualization, Methodology, Investigation, Writing – original draft, Writing – review & editing, Supervision. **Hinrich Kaiser:** Conceptualization, Methodology, Investigation, Writing – original draft, Writing – review & editing, Supervision, Project administration, Funding acquisition.

#### Declaration of Competing Interest

The authors declare that there is no conflicts of interest.

#### Acknowledgments

For their assistance with fieldwork, we thank the students of the Winter 2012, Summer 2012, and Summer 2013 surveys, Zach Brown, Kevin Burns, Melissa Carrillo, Jester Ceballos, Britta Döring, Scott Heacox, Stephanie Hughes, Andrew Kathriner, Eric Leatham, Aaren Marsh, Gloria Morales, John Paris, Julia Pozo, Justin Rader, Claudia Rivas, Caitlin Sanchez, Dan Suzio, David Taylor, Marianna Tucci, Franziska Wagner, and MJ Weil. Our fieldwork gained significantly in terms of biological productivity and cultural learning due to the hard work of our young Timorese colleagues, Venancio Lopes Carvalho, Luis Lemos de Araujo, Agivedo Varela Ribeiro, and Zito Afranio Soares. We also thank our colleague Sven Mecke, who not only worked with us during several phases of fieldwork but who became a full-fledged professional as we were watching. We acknowledge the help of Alberto A. Guglielmone for recommending several publications during our literature survey. Our studies in Timor-Leste were made possible only through the interest and gracious support of three government leaders, President and Nobel Peace Laureate José Ramos-Horta, former Prime Minister Xanana Gusmão, and former Minister for the Council of Ministers Ágio Pereira. Their friendship to our cause has been appreciated very much. We also wish to express our gratitude to Claudia Abate-Debat, former special advisor in the Prime Minister's Office, whose behind-the-scenes activities on our behalf made our goals come to fruition. For issuing collecting and export permits, we thank Manuel Mendes, Director of National Parks. Our research in Timor-Leste supported financially by student travel grants from the Associated Student Body at Victor Valley College and the Victor Valley College Foundation. This paper is Contribution No. 26 from the Tropical Research Initiative at Victor Valley College. Fábrício H. Oda received a postdoctoral fellowship from the Fundação Cearense de Apoio ao Desenvolvimento Científico e

Tecnológico (Grant n. 6471073/2017). Thiago F. Martins is supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP process n. 2019/03167-0, 2020/05987-1).

#### References

- Anastos, G., 1950. The scutate ticks, or Ixodidae, of Indonesia. Entomol. Am. 30, 1–144.
- ASIH [American Society of Ichthyologists and Herpetologists], 2004. Guidelines for the Use of Live Amphibians and Reptiles in Field and Laboratory Research, Second Edition. Animal Care Committee, ASIH, Athens, p. 43.
- Audy, J.R., Nadchatram, M., Boo-Liat, L., 1960. Malaysian parasites XLIX. Host distribution of Malayan ticks (Ixodoidea). Stud. Inst. Med. Res. Fed. Malaya 29, 225–246.
- Auffenberg, T., 1988. *Amblyomma helvolum* (Acarina: Ixodidae) as a parasite of varanid and scincid reptiles in the Philippines. Int. J. Parasitol. 18, 937–945.
- Burridge, M.J., Simmons, L.A., 2003. Exotic ticks introduced into the United States on imported reptiles from 1962 to 2001 and their potential roles in international dissemination of diseases. Vet. Parasitol. 113, 289–320.
- Burridge, M.J., 2011. Non-native and invasive ticks. Threats to human and animal health in the United States. University of Florida Press, Gainesville.
- Chao, L.-L., Hsieh, C.-K., Shih, C.-M., 2013. First report of *Amblyomma helvolum* (Acar: Ixodidae) from the Taiwan stink snake, *Elaphe carinata* (Reptilia: Colubridae), collected in southern Taiwan. Ticks Tick-Borne Dis. 4, 246–250.
- Clark, L.G., Doten, E.H., 1995. Ticks on imported reptiles into Miami International Airport: November 1994 through January 1995. Proceedings for the Veterinary Epidemiology and Economics Symposium. United States Department of Agriculture, Fort Collins, pp. 1A17–1A25.
- Chen, Z., Yang, X., Bu, F., Yang, X., Yang, X., Liu, J., 2010. Ticks (Acar: Ixodoidea: Argasidae, Ixodidae) of China. Exp. Appl. Acarol. 51, 393–404.
- Dias, J.A.T.S., 1988. A contribution towards the knowledge of ticks in the Timor Island. Series: Zoologia 15, 89–96.
- Doornbos, K., Sumrandee, C., Ruang-Areerate, T., Baimai, V., Trinachartvanit, W., Ahantaraig, A., 2013. *Rickettsia* sp. closely related to *Rickettsia rauoultii* (Rickettsiales: Rickettsiaceae) in an *Amblyomma helvolum* (Acarina: Ixodidae) tick from a *Varanus salvator* (Squamata: Varanidae) in Thailand. J. Med. Entomol. 50, 217–220.
- Durden, L.A., Merker, S., Beati, L., 2008. The tick fauna of Sulawesi, Indonesia (Acar: Ixodoidea: Argasidae and Ixodidae). Exp. Appl. Acarol. 45, 85–110.
- GBIF.org, 2020. GBIF Home Page. <https://www.gbif.org/> (accessed 25 July 2020).
- Ghosh, H.S., Bansal, G.C., Gupta, S.C., Ray, D., Khan, M.Q., Irshad, H., Shahiduzzaman, M., Seitzer, U., Ahmed, J.S., 2007. Status of tick distribution in Bangladesh, India and Pakistan. Parasitol. Res. 101 (2), 207–216 (Suppl).
- Ghosh, H.S., Roy, S., Sanyal, A.K., Misra, K.K., 2017. Microscopic anatomy of the Haller's organ of snake ticks. Int. J. Acarol. 43, 1–9.
- Ghosh, H.S., Sanyal, A.K., Misra, K.K., 2018. Studies on the surface ultrastructure of a snake tick, *Amblyomma helvolum* (Acar: Ixodidae). J. Anim. Sci. Vet. Med. 3, 75–89.
- Ghosh, H.S., Sanyal, A.K., Misra, K.K., 2021. Survey for snake-tick (Ixodida) association in some districts of West Bengal, India. Persian J. Acarol. 10, 55–67.
- Guerrero Sanchez, S., 2019. Ecology and Health of the Asian Water Monitor Lizard (*Varanus salvator*) in the Fragmented Landscape of the Kinabatangan Floodplain, Sabah, Malaysia. Cardiff University, Cardiff. PhD Thesis.
- Guglielmone, A.A., Robbins, R.G., Apanaskevich, D.A., Petney, T.N., Estrada-Peña, A., Horak, I.G., 2014. The hard ticks of the world: (Acar: Ixodida: Ixodidae). Springer, Dordrecht, Heidelberg, New York, London.
- Guglielmone, A.A., Sánchez, M.E., Franco, L.G., Nava, S., Rueda, L.M., Robbins, R.G., 2015. Names of species of hard ticks. Instituto Nacional de Tecnología Agropecuaria, Rafaela, Argentina. <http://rafaela.inta.gob.ar/nombresgarrapatas>. (accessed 26 March 2022).
- Guglielmone, A.A., Robbins, R.G., 2018. Hard Ticks (Acar: Ixodida: Ixodidae) Parasitizing Humans: A Global Overview. Springer, Dordrecht, Heidelberg, New York, London.
- Hirst, S., Hirst, L.F., 1910. XXXV. Descriptions of five new species of ticks (Ixodidae). Ann. Mag. Nat. Hist. 6, 299–308.
- Hoogstraal, H., Santana, F.J., Van Peenen, P.D., 1968. Ticks (Ixodoidea) of Mt. Sontra, Danang, Republic of Vietnam. Ann. Entomol. Soc. Am. 61, 722–729.
- Kaenkan, W., Nooma, W., Chelong, I., Baimai, V., Trinachartvanit, W., Ahantaraig, A., 2020. Reptile-associated *Borrelia* spp. in *Amblyomma* ticks, Thailand. Ticks Tick-Borne Dis. 11, 101315.
- Kaiser, H., Carvalho, V.L., Ceballos, J., Freed, F., Heacox, S., Lester, B., Richards, S.J., Trainor, C.R., Sanchez, C., O'Shea, M., 2011. The herpetofauna of Timor-Leste: a first report. ZooKeys 109, 19–86.
- Kaiser, H., Sanchez, C., Heacox, S., Kathriner, A., Varela Ribeiro, A., Afranio Soares, Z., Lemos de Araujo, L., Mecke, S., O'Shea, M., 2013. First report on the herpetofauna of Ataúro Island, Timor-Leste. Check List 9, 752–762.
- Keirans, J.E., 1984. George Henry Falkiner Nuttall and the Nuttall Tick Catalogue. U. S. Department of Agriculture Miscellaneous Publications, Washington.
- Keirans, J.E., Durden, L.A., 2001. Invasion: exotic ticks (Acar: Argasidae, Ixodidae) imported into the United States. A review and new records. J. Med. Entomol. 38, 850–861.
- Kho, K.L., Koh, F.X., Tay, S.T., 2015. Molecular evidence of potential novel spotted fever group rickettsiae, *Anaplasma* and *Ehrlichia* species in *Amblyomma* ticks parasitizing wild snakes. Parasites Vectors 8, 112.
- Kho, K.L., Tan, P.E., Tay, S.T., 2019. Diversity of rickettsiae in feeding and questing ticks collected from a malaysian forest reserve area. J. Med. Entomol. 56, 547–552.

- Kim, N., 2020. A reticulated python infested with the tick, *Amblyomma helvolum*. Singapore Biodivers. Rec. 2020, 68.
- King, D., Green, B., 1999. Monitors: The Biology of Varanid Lizards. Krieger Publishing Company, Malabar, Florida, USA.
- Kohls, G.M., 1950. Ticks (Ixodoidea) of the Philippines. Nat. Inst. Health Bull. 192, 28.
- Kohls, G.M., 1957. Malaysian parasites. XVIII. Ticks (Ixodoidea) of Borneo and Malaya. Stud. Inst. Med. Res. Malaya 28, 65–94.
- Kolonin, G.V., 1995. Review of the ixodid tick fauna (Acar: Ixodidae) of Vietnam. J. Med. Entomol. 32, 276–282.
- Krijgsman, B.J., Ponto, S.A.S., 1931. Die Verbreitung der Zecken in Niederländisch-Ostindien. Ztschr. Parasitenk. 4, 140–146.
- Kwak, M.L., 2020. A checklist and key to the tick fauna (Acar: Ixodidae, Argasidae) of Pulau Tioman, Malaysia. Exp. Appl. Acarol. 81, 51–58.
- Lazell, J.D., Keirans, J.E., Samuelson, G.A., 1991. The Sulawesi black racer, *Coluber (Ptyas) dipsas*, and remarkable ectoparasite aggregation. Pac. Sci. 45, 355–361.
- Livezey, R.L., 1958. Procaine hydrochloride as a killing agent for reptiles and amphibians. Herpetologica 13, 280.
- Mao, J.-J., Norval, G., Robbins, R.G., Wong, S.T., 2019. An instance of *Boiga dendrophila dendrophila* (Boie, 1827) (Reptilia: Colubridae) being parasitized by *Amblyomma helvolum* Koch, 1844 (Acar: Ixodidae), with comments about the attachment sites of this tick species. Acarologia 59, 115–119.
- Monk, K.A., de Fretes, Y., Reksodiharjo-Lilley, G., 1997. The Ecology of Nusa Tenggara and Maluku. The Ecology of Indonesia Series, Volume V. Periplus Editions, Singapore.
- Nadchatram, M., Domrow, R., Ng, C.K., 1966. Parasitic Acarina of the mammals. Bull. Natl. Mus. State Singapore 34, 129–140.
- Nakatsui, K., 1937. On the tick parasited on the lizard and snake from Philippine Is. [sic] Acta Arachnol. 2, 65–67.
- Neumann, L.G., 1899. Révision de la famille des ixodidés (3<sup>e</sup> mémoire). Mém. Soc. Zool. Fr., Paris 12, 107–294.
- Neumann, L.G., 1901. Révision de la famille des ixodidés (4<sup>e</sup> mémoire). Mém. Soc. Zool. Fr., Paris 14, 249–372.
- Neumann, L.G., 1910. Sur quelques espèces d'Ixodidae nouvelles ou insuffisamment connues. Annal. Sci. Nat. Zool. Sér. 9, 161–176.
- Neumann, L.G., 1911. Ixodidae. Das Tierreich 26, 1–169.
- O'Shea, M., Sanchez, C., Heacox, S., Kathriner, A., Lopes Carvalho, V., Varela Ribeiro, A., Afranio Soares, Z., Lemos de Araujo, L., Kaiser, H., 2012. First update to herpetofaunal records from Timor-Leste. Asian Herpetol. Res. 3, 114–126.
- O'Shea, M., Sanchez, C., Kathriner, A., Mecke, S., Lopes Carvalho, V., Varela Ribeiro, A., Afranio Soares, Z., Lemos de Araujo, L., Kaiser, H., 2015. Herpetological diversity of Timor-Leste: updates and a review of species distributions. Asian Herpetol. Res. 6, 73–131.
- Oudemans, A.C., 1928. Acarologische Aanteekeningen XCI. Entomol. Ber. VII 317–328.
- Parrish, D.W., 1971. The Occurrence, Geographical Distribution and Wild Vertebrate Host Relationships of Ticks (Ixodoidea) on Luzon Island, Philippines, with Descriptions of Three New Species. Oklahoma State University, Stillwater. Ph.D. dissertation.
- Petney, T.N., Keirans, J.E., 1995. Ticks of the genera *Amblyomma* and *Hyalomma* (Acar: Ixodidae) in South-east Asia. Trop. Biomed. 12, 45–56.
- Pianka, E.R., King, D.R., King, R.A., 2004. Varanoid Lizards of the World. Indiana Univ. Press, Bloomington, Indiana, USA xiii + 588 pp.
- Robinson, L.E., 1926. Ticks. A monograph of the Ixodoidea. Part IV. The genus *Amblyomma*. Cambridge University Press, Cambridge.
- Sanchez, C., Lopes Carvalho, V., Kathriner, A., O'Shea, M., Kaiser, H., 2012. First report on the herpetofauna of the Oecusse District, an exclave of Timor-Leste. Herpetol. Notes 5, 137–149.
- Sanyal, A.K., De, S.K., 1992. Ixodid ticks (Acar: Ixodidae), in: Ghosh, A.K. (Ed.), State Fauna Series 3: Fauna of West Bengal, Part - 3 (Arachnida and Acari). Zoological Survey of India, Calcutta, pp.17–60.
- Schulze, F., 1933. Ixodidae der deutschen limnologischen Sunda-Expedition. Archiv Hydrobiol. Suppl. Bd. XII, "Tropische Binnengewässer, Band IV" 490–503.
- Shea, G.M., 2012. On the identity of the type species of *Sphenomorphus* (Squamata: Scincidae): *Lygosoma melanopogon* Duméril and Bibron 1839 and a note on a new scalation character of the pes in *Sphenomorphus*. Zootaxa 3490, 1–29.
- Sharif, M., 1928. A revision of the Indian Ixodidae with special reference to the collection in the Indian Museum. Rec. Indian Mus. 30, 217–344.
- Simmons, L.-A., Stadler, C.K., Burridge, M.J., 2002. Introduction of the exotic tick *Amblyomma helvolum* Koch (Acar: Ixodidae) into the United States on imported cobras (Squamata: Elapidae). Int. J. Acarol. 28, 45–48.
- Sumrandee, C., Baimai, V., Trinachartvanit, W., Ahantarig, A., 2015. *Hepatozoon* and *Theileria* species detected in ticks collected from mammals and snakes in Thailand. Ticks Tick-borne Dis 6, 309–315.
- Sumrandee, C., Hirunkanokpun, S., Doornbos, K., Kitthawee, S., Baimai, V., Grubhoffer, L., Trinachartvanit, W., Ahantarig, A., 2014b. Molecular detection of Rickettsia species in *Amblyomma* ticks collected from snakes in Thailand. Ticks Tick-borne Dis 5, 632–640.
- Sumrandee, C., Hirunkanokpun, S., Grubhoffer, L., Baimai, V., Trinachartvanit, W., Ahantarig, A., 2014a. Phylogenetic relationships of *Francisella*-like endosymbionts detected in two species of *Amblyomma* from snakes in Thailand. Ticks Tick-borne Dis 5, 29–32.
- Tabug, M.A., Necesito, L.V., 2021. *Lycodon capucinus* (Common Wolf Snake). Ectoparasitism. Herp. Rev. 52, 426–427.
- Tanskul, P., Stark, H.E., Inlao, I., 1983. A checklist of ticks of Thailand (Acar: Metastigmata: Ixodoidea). J. Med. Entomol. 20, 330–341.
- Uetz, P., Freed, P., Aguilar, R., Hošek, J., 2021. The reptile database. <http://www.reptile-database.org> (accessed 21 November 2021).
- USDA, 1971. National Tick Surveillance Program, Calendar Year 1970. Animal Health Division, Agricultural Research Service. US Department of Agriculture, Hyattsville, MD.
- Voltzit, O.V., Keirans, J.E., 2002. A review of Asian *Amblyomma* species (Acar, Ixodida, Ixodidae). Acarina 10, 95–136.
- Warburton, C., 1910. On two collections of Indian ticks. Parasitology 3, 395–407.
- Warburton, C., 1926. Fauna Simalurensis. Ixodidae. Treubia 8, 3–4.